

# Synthesis, Characterization, and Application of Molecularly Imprinted Polymer - Modified Silica Gel for Andrographolide Purification from *Andrographis paniculata* Burm.F.Ness Methanol Extract

*by Diki Prayugo Wibowo*

---

**Submission date:** 20-Jul-2023 09:19PM (UTC-0700)

**Submission ID:** 2134384899

**File name:** D2-TJNPR-2023-M323\_Reviewed\_1\_exclude\_biblio.pdf (305.59K)

**Word count:** 2086

**Character count:** 11716

# Synthesis, Characterization, and Application of <sup>3</sup> Molecularly Imprinted Polymer - Modified Silica Gel for Andrographolide Purification from *Andrographis paniculata* *Burm.F.Ness* Methanol Extract

## Abstract

**Molecularly Imprinted Polymer** -Modified silica is silica gel coated with Molecularly Imprinted Polymer (MIP) which has cavities with the same shape, size, and functional groups as the template molecule, allowing for selective absorption. Therefore, this study aimed to synthesize, characterize, and apply MIP-modified silica for purification of andrographolide from the *Andrographis paniculata* *Burm.F.Ness* methanol extract. To synthesize the MIP-modified silica, the surface imprinting method was employed, with APTES, TEOS, acetic acid, and andrographolide, as the functional monomer, cross-linking agent, catalyst, and template molecule, respectively. Modified silica was characterized using FTIR and SEM. Its performance was tested by determining the imprinting factor. The purity of purified methanol extract was tested using HPLC. FTIR and SEM characterization confirmed the presence of MIP polymer on the surface of silica gel. Its imprinting factor was 1.22. Purification results of the *Andrographis paniculata* *Burm.F.Ness* extract demonstrated that modified silica improved the purity of methanol extract from 1.46 % to 65%.

**Keywords:** Modified Silica Gel, Andrographolide, Surface Imprinting, Silica gels

## Introduction

Andrographolide, the primary compound discovered in the *Andrographis paniculata* *Burm.F.Ness* leaves, exhibits various biological activities, including anti-inflammatory, antidiabetic, antiplatelet, antibacterial, antiviral, and immunostimulant effects<sup>1</sup>. The pharmacological activity of andrographolide and the growing interest in natural treatments promote extensive study on its isolation from the *Andrographis paniculata* *Burm.F.Ness* plant. This compound relies on recrystallization and *chromatographic* methods for its purification<sup>2</sup>. However, multiple recrystallization processes are required to obtain a high-purity isolate, while chromatography employs environmentally hazardous organic solvents. Therefore, there is a need to develop purification method that is simpler and more environmentally friendly. <sup>12</sup> Molecularly Imprinted Polymer (MIP), which was coated on silica gel in this study, is an alternative method for andrographolide purification<sup>3</sup>.

Modified silica refers to silica gel coated with MIP which is a polymer with specific and selective cavities designed to imprint the target or template molecule, in terms of its shape, size,

and functional groups<sup>4</sup>. It comprises molecules or compounds to be imprinted (template), functional monomers, crosslinkers, initiators, and porogen solvents<sup>4</sup>.

The coating of silica with MIP aims to prevent the expansion of the polymer structure, which could alter the imprint of the target compound due to interactions with organic solvents. This modification enhances the specificity and selectivity of silica gel as an effective adsorbent by modifying silanol groups<sup>5</sup>. A previous study conducted by Yang et al.<sup>6</sup> on the Bisphenol A compound, synthesized MIP-modified silica, resulting in a recovery percentage of 99.43% which was significantly higher than the conventional isolation method of 3.6%.

The synthesized modified silica was characterized using Scanning Electron Microscope (SEM) and Fourier Transform Infrared (FTIR) techniques. The results showed that it can be utilized as an adsorbent for Solid Phase Extraction (SPE).

## Methods

### Tools

The study involved the use of several tools, including an analytical balance (Ohaus®), a set of reflux and distillation apparatus, an orbital shaker (IKA® KS 130 Basic), an ultrasonicator (Elmasonic® S 30H), a centrifuge (Hettich® EBA 20), a UV-Visible spectrophotometer (Shimadzu® UV 1800), an SEM (Hitachi SU 3500®), an FTIR (Thermo® Scientific), a micropipette (FISHER Elite®), and glassware commonly used in the Chemistry Laboratory.

### Materials

The materials used in this study were silica gel 60 (70-230 mesh ASTM) for column chromatography (MERCK®), andrographolide (Tokyo Chemical Industry®), (3-aminopropyl) triethoxysilane (APTES) (Tokyo Chemical Industry®), tetraethoxysilane (TEOS) (MERCK®), methanol pro-HPLC (MERCK®), glacial acetic acid (Fulltime®), and demineralized water (Amidis®), all the reagent used were analytical grade.

### MIP-Modified Silica Gel Synthesis

Before application, silica gel was activated by refluxing it for 12 hours using a 3 M HCl solution with a volume of 200 mL. Subsequently, it was filtered and washed with demineralized water to neutralize silica. The final product was then dried in an oven at 110°C for 12 hours and stored in a closed container, as described by<sup>7</sup>.

3-aminopropyl triethoxysilane (APTES) as a functional monomer and andrographolide solution (template) in methanol were mixed using a magnetic stirrer at room temperature for 30 minutes. Furthermore, 1 g of activated silica gel was added to the mixture, along with

tetraethoxysilane (TEOS) as a crosslinker and glacial acetic acid. The mixing process was continued for 24 hours. Subsequently, modified silica was filtered and the template was removed using methanol: acetic acid solvent at 9:1 through sonication for 7 hours. The sonication was continued until the absorption wavelength of andrographolide in extracting solvent was no longer detected by UV-Vis spectrophotometry. Finally, the synthesized modified silica was dried in an oven at 100°C for 8 hours, following the method described by 7, 8.

The composition of the components constituting the modified MIP-silica gel can be seen in the table 1.

### Characterization of MIP -Modified Silica Gel using SEM

The surface morphology of modified silica was observed using SEM. For the analysis, the sample was placed on a double-sided metal plate, and subsequently, it was gold-plated under vacuum conditions. The scanning process was conducted using a current of 60 mA and an electric power of 15 V.

### Characterization of MIP - Modified Silica using FTIR

To analyse the functional groups in modified silica, FTIR spectroscopy was employed. Approximately 50 mg of the sample was placed in ZnSe ATR and IR radiation was performed at wave numbers that ranged from 400 to 4000 cm<sup>-1</sup>.

### Imprinting Factor Determination

Imprinting factor was determined using equations 1 and 2:

$$KD = \frac{(C_i - C_f)V}{C_f W} \quad 1$$

$$IF = \frac{KDMS}{KDUMS} \quad 2$$

KD is the distribution coefficient, C<sub>i</sub> and C<sub>f</sub> (mg/L) are andrographolide concentration before and after the adsorption experiment, V (L) is the volume of andrographolide solution, and W (g) is the polymer weight. IF stands for imprinting factor, KD Ms is the distribution coefficient of modified silica, and KD UMS is the distribution coefficient of modified silica without the addition of template compounds<sup>9-13</sup>.

### Application of modified silica for andrographolide purification

A total of 5 grams of modified silica was mixed with 100 mL of the *Andrographis paniculata* Burm.F.Nes methanol extract solution using a shaker. The mixing process lasted for 1 hour at a speed of 240 rpm. Upon completion, the mixture was separated into filtrate and modified silica was separated. Furthermore, the solvent was evaporated, and purification results were characterized using UV-Vis and FTIR spectrophotometry. Finally, the purity of the purified isolate was determined using HPLC<sup>8</sup>.

## RESULTS AND DISCUSSION

### Modified Silica Gel Synthesis

During the activation process of silica gel, there exists a change from SiO<sub>2</sub> to Si-OH, as shown in **Figure 1**. This enables it to readily form polymers with TEOS. The success of the activation process was confirmed by increased absorption within the wave numbers of 3000 to 3500 cm<sup>-1</sup>, indicating that the hydroxyl groups in the activated silica gel had increased. Synthesis process of modified silica was described using the following scheme.

### Characterization of Modified Silica using FTIR and SEM

#### Characterization Results Using SEM

The SEM characterization results provided information about the surface morphology of modified silica gel. **Figure 2** shows the difference in the surface morphology between silica gel and modified silica gel. Modified silica gel exhibited a rougher texture. Additionally, a change in particle size was observed in silica gel before and after modification. This occurred due to the formation of MIP on its surface.

#### Characterization Results using FTIR

The objective of FTIR characterization of modified silica was to observe the formation of functional groups, following the synthesis process. The surface modification of silica gel involved the hydrolysis and conjugation of TEOS and APTES, leading to the elimination of alkyl groups in both compounds. This was indicated by the absence of absorption at wavenumbers 2980 and 2670 cm<sup>-1</sup> as well as 2980 and 2850 cm<sup>-1</sup>. Furthermore, the presence of absorption within the wavenumber range of 1000-1100 cm<sup>-1</sup> showed the successful formation of MIP on the silica gel surface<sup>14</sup>, attributed to the presence of Si-O-Si groups<sup>14</sup>.

FTIR characterization results of modified silica did not provide a clear overview of the interactions between andrographolide and the polymer. However, they could indicate the success of synthesis process through the absence of absorption at wavenumbers 2980 and 2670 cm<sup>-1</sup> as well as 2980 and 2850 cm<sup>-1</sup>, as shown in **Figure 3**. The differences in the spectrum shape

between modified silica and its constituent materials (silica gel, TEOS, and APTES) suggested the occurrence of a polymerization process<sup>14</sup>.

### **Imprinting Factor determination**

Imprinting factor is a parameter that shows the performance of MIP on modified silica gel, the greater the imprinting factor, the better the MIP of modified silica gel performance. MIP – modified silica gel showed good performance since it has an IF value more than 1<sup>8,9</sup>.

The composition of a molecularly imprinted polymer (MIP) coating on silica gel significantly influences the imprinting factor. Consequently, variations in the imprinting factor values are observed between MIP- silica gel fabricated using a functional monomer ratio of 1:4 (MSA) and those prepared with a template: functional monomer ratio of 1:5 (MSB)<sup>15</sup>. The results of the imprinting factor test can be observed in [Table 1](#).

### **Modified Silica Gel Application for Andrographolide Purification**

The purification process involved mixing the Sambiloto methanol extract with a certain amount of modified silica gel at the optimal contact time. The modified MIP-silica gel used for purification was the MIP-silica gel modification with the best imprinting factor (MSA). The purified extract showed an increase in purity from 1.46 % to 65.97% (retention time: 3.520). This was higher compared to purification with unmodified silica gel from 1.46 % to 20.57%. (RT: 3.352). The chromatograms of methanol extract before and after purification are shown in Figure 4.

### **CONCLUSION**

Based on characterization of modified silica gel using SEM, FTIR, and imprinting factor test, MIP was formed on the surface of silica gel, which is selective for andrographolide. The increased selectivity, achieved through modification with MIP, was further evidenced by the higher purity obtained in the isolate purified using modified silica gel. Therefore, MIP-modified silica gel proved to be a valuable tool for purification process of andrographolide from the *Andrographis paniculata* *Burm.F.Nes* methanol extract.

### **Conflict of interest**

All the authors affirm that they have no conflicts of interest in relation to this research.


### **Authors' Declaration**

The authors affirm that the content presented in this article is entirely original and take full responsibility for any claims arising from its content.


**Acknowledgements**

We would like to express our sincere appreciation and gratitude to Sekolah Tinggi Farmasi Indonesia (STFI) Bandung for their invaluable support in providing laboratory facilities for this research project.

**References (for appropriate referencing style consult journal author guideline)**




**Figure 1. Illustration of Modified silica gel synthesis. a : template removal process; b : rebinding of target molecules to the MIP cavity ; c : modified silica gel before template removal, d: modified silica gel after template removal; e : silica gel before modification**




**Figure 2. Characterization results using SEM, a: silica gel b: MIP - modified silica gel**





**Figure 3: Characterization results using FTIR. a: activated silica gel; b: APTES; c: TEOS; d : MSA ; e : MSB**



**Figure 4: The chromatogram of the purity test results of the methanol extract of *Andrographis paniculata* before purification (b), after purification using silica gel (d), and after purification using modified MIP-silica gel (c), were compared with the standard (a).**

**Images should be of high quality**

**Table 1: Composition of modified MIP-silica gel**

<b>Adsorbent</b>	<b>Materials</b>	<b>Total</b>
Modified Silica A (MSA)	andrographolide	280 mg
	APTES	0.94 mL
	TEOS	18 mL
	Acetic acid glacial	1 mL
	silica gel	1 g
	Methanol	30 mL
MSA without template addition (NSA)	andrographolide	0
	APTES	0.94 mL
	TEOS	18 mL
	Acetic acid glacial	1 mL
	silica gel	1 g
	Methanol	30 mL
Modified Silica B (MSB)	andrographolide	280 mg
	APTES	1.175 mL
	TEOS	4.43 mL
	Acetic acid glacial	1 mL
	silica gel	1 g
	Methanol	30 mL
MSB without template addition (NSB)	andrographolide	0
	APTES	0.24 mL
	TEOS	4.43 mL
	Acetic acid glacial	1 mL
	silica gel	1g
	Methanol	30 mL

Note : MSA and MSB had different comparison of template molecule and functional monomer. The comparison of template molecules : functional monomer of MSA 1:4. While the comparison in MSB was 1:5.

**Table 2: The result of Imprinting Factor determination**

<b>Adsorbent</b>	<b>KD</b>	<b>IF</b>
------------------	-----------	-----------

---

MSA	0.104075	
NSA	0.085477	1.22
MSB	0.084338	
NSB	0.085477	0.99

Note : KD is Distribution coefficient of the sample, IF is imprinting factor of the sample.

# Synthesis, Characterization, and Application of Molecularly Imprinted Polymer - Modified Silica Gel for Andrographolide Purification from *Andrographis paniculata* Burm.F.Ness Methanol Extract

## ORIGINALITY REPORT

13%

SIMILARITY INDEX

4%

INTERNET SOURCES

11%

PUBLICATIONS

3%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Wiwin Winingsih, Slamet Ibrahim, Sophi Damayanti. "Purification of Andrographolide Methanolic Extract Using Molecularly Imprinted Polymer Prepared by Precipitation Polymerization", *Scientia Pharmaceutica*, 2022  
Publication 2%
- 2 Li, Min-ting, Yong-yan Zhu, Li Li, Wen-na Wang, Yong-guan Yin, and Quan-hong Zhu. "Molecularly Imprinted Polymers on a Silica Surface for the Adsorption of Tobacco-specific Nitrosamines in Mainstream Cigarette Smoke", *Journal of Separation Science*, 2015.  
Publication 1%
- 3 Mingkun Gao, Yuhang Gao, Ge Chen, Xiaodong Huang, Xiaomin Xu, Jun Lv, Jing Wang, Donghui Xu, Guangyang Liu. "Recent Advances and Future Trends in the Detection of Contaminants by Molecularly Imprinted 1%

## Polymers in Food Samples", Frontiers in Chemistry, 2020

Publication

---

4

Mao Suganami, So Konno, Ryo Maruhashi, Daisuke Takagi et al. "Expression of flavodiiron protein rescues defects in electron transport around PSI resulting from overproduction of Rubisco activase in rice", Journal of Experimental Botany, 2022

Publication

---

1 %

5

E. Díez, J. M. Gómez, A. Rodríguez, I. Bernabé, J. Galán. "Recovery of Gallium from Aqueous Solution through Preconcentration by Adsorption/Desorption on Disordered Mesoporous Carbon", Journal of Sustainable Metallurgy, 2021

Publication

---

1 %

6

Akl, M.A.A.. "Organically modified silica gel and flame atomic absorption spectrometry: employment for separation and preconcentration of nine trace heavy metals for their determination in natural aqueous systems", Microchemical Journal, 200410

Publication

---

1 %

7

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)

Internet Source

---

1 %

8

[www.science.gov](http://www.science.gov)

Internet Source

---

1 %

9	Submitted to Liverpool John Moores University Student Paper	1 %
10	academicjournals.org Internet Source	1 %
11	Hitoshi Shiku, Isamu Uchida, Tomokazu Matsue. "Microfabrication of Alkylsilanized Glass Substrate by Electrogenenerated Hydroxyl Radical Using Scanning Electrochemical Microscopy", Langmuir, 1997 Publication	<1 %
12	Marziyeh Poshteh Shirani, Behzad Rezaei, Ali A. Ensafi, Mohammad Ramezani. "Development of an eco-friendly fluorescence nanosensor based on molecularly imprinted polymer on silica-carbon quantum dot for the rapid indoxacarb detection", Food Chemistry, 2020 Publication	<1 %
13	Peter Spégel. "Molecularly imprinted microparticles for capillary electrochromatography: Studies on microparticle synthesis and electrolyte composition", Electrophoresis, 09/2001 Publication	<1 %
14	adtplus.arcanum.hu Internet Source	<1 %

15

[journal.unhas.ac.id](http://journal.unhas.ac.id)

Internet Source

&lt;1 %

16

Goswami, A.. "1,8-Dihydroxyanthraquinone anchored on silica gel: synthesis and application as solid phase extractant for lead(II), zinc(II) and cadmium(II) prior to their determination by flame atomic absorption spectrometry", *Talanta*, 20021016

Publication

&lt;1 %

17

[www.researchgate.net](http://www.researchgate.net)

Internet Source

&lt;1 %

18

Andac, M.. "Ion-imprinted beads for molecular recognition based mercury removal from human serum", *International Journal of Biological Macromolecules*, 20070130

Publication

&lt;1 %

19

Chaiyavat Chaiyasut. "Estimation of the dissociation constants for functional groups on modified and unmodified silica gel supports from the relationship between electroosmotic flow velocity and pH", *Electrophoresis*, 05/2001

Publication

&lt;1 %

20

Benoit Guieysse, Mathieu Le Noir, Bo Mattiasson. "Removal of Endocrine Disrupting Compounds Using Molecularly Imprinted

&lt;1 %

# Polymers: A Review", American Chemical Society (ACS), 2010

Publication

---

21

Zahra Mohammadbagheri, Abbas Rahmati, Parisa Hoshyarmanesh. "Synthesis of a novel superabsorbent with slow-release urea fertilizer using modified cellulose as a grafting agent and flexible copolymer", International Journal of Biological Macromolecules, 2021

Publication

---

<1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off



# Synthesis, Characterization, and Application of Molecularly Imprinted Polymer - Modified Silica Gel for Andrographolide Purification from *Andrographis paniculata* Burm.F.Ness Methanol Extract

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---

PAGE 10

---