

# Hesti Riasari\_PCPR

*by* Hesti Riasari

---

**Submission date:** 01-Feb-2023 10:20AM (UTC+0800)

**Submission ID:** 2003761043

**File name:** Hesti\_Riasari\_PCPR.pdf (242.6K)

**Word count:** 1963

**Character count:** 10745

## Anti-Inflammatory Activity of *Artocarpus altilis* (Parkinson) Fosberg in Wistar Male Rats

Hesti Riasari, Sani Nurlaela, Ginanjar C. Gumilang  
Indonesia School of Pharmacy, Bandung, West Java, Indonesia

### Abstract

Inflammation is a local reaction in vascular tissue characterized by several symptoms, e.g., swelling, redness, and pain. Indonesia is one of the countries with high biodiversity, including traditional medicinal plants. *Artocarpus altilis* is one of the plants that is widely used to treat inflammation. There is limited information on biological activity of *A. altilis*. This study was performed to evaluate anti-inflammatory properties of *A. altilis* detached and attached leaves in Wistar male rats. *A. altilis* was extracted using soxhlet method at 67.4 °C with methanol solvent. Inflammation was induced by the administration of carageenan to the rats paw. Subsequently, the extract of *A. altilis* were orally administered. The edema were measured using plethysmometer for 6 hours. We found that there were differences in anti-inflammatory activity between detached and attached leaves. The inhibition of edema in attached leaves were 50% and 53.33% for the concentration of 50 mg and 100 mg, respectively. The greater inhibition was observed in the detached leaves, with 73.33% and 76.67% inhibition, for the concentration of 50 mg and 100 mg, respectively. Nevertheless, the inhibition percentage was still below diclofenac sodium as a positive control (83.33%). In conclusion, *A. altilis* leaves extracts showed good anti-inflammatory properties and has the potential for development of anti-inflammatory drug.

**Keywords:** *Artocarpus altilis*, carageenan, inflammation, edema

### Introduction

Inflammation is a local response to cellular injury, characterized by capillary dilatation, swelling, redness, and pain. The treatment includes the administration of anti-inflammatory drugs. Nevertheless, many side effects were associated with the use of these drugs, such as gastrointestinal ulcer, bleeding, kidney failure, etc.<sup>1-3</sup>

<sup>1</sup> The increasing use of herbal medicine and phytonutrients has been observed during the last decade. The use of medicinal plant is preferred by many individuals since it may have fewer side effects compared to the conventional treatment. It is estimated that up to four billion people worldwide rely on herbal medicinal products.<sup>4,5</sup>

---

Corresponding author: Hesti Riasari. Indonesia School of Pharmacy, Bandung, West Java, Indonesia. Email: hmm\_riasari@yahoo.com

Received: 13 March 2019. Revised: 6 April 2019. Published: 25 April 2019.

**Table 1. Characteristics of *A. Altilis* Leaves**

Leaves	Ash (%)	Water Soluble Compounds (%)	Ethanol Soluble Compounds (%)
Attached	26.00	32.94	8.92
Detached	27.36	35.94	12.44

Indonesia is one of the country with highest biodiversity, including traditional medicinal plants.<sup>6</sup> *A. altilis* is one of the Indonesian traditional medicinal plants that is used to treat inflammation. Previous study showed that detached and attached leaves had significant differences in defence response.<sup>7</sup> Two types of leaves might have different metabolites which can influence its biological activities. Nevertheless, there is limited information on biological activity of the variety of *A. altilis* leaves. This study was performed to evaluate anti-inflammatory properties of *A. altilis* in Wistar male rats.

## Methods

### Materials

The materials used in this study included the leaves of *A. altilis* (Park.) Fosberg which were obtained from the Cipamokolan, Bandung, West Java, Indonesia, methanol, distilled water, carrageenan, diclofenac sodium, and several phytochemical screening reagents, e.g., ammonia, chloroform, hydrochloride, gelatin solution, amyl alcohol, ether, vanilla, H<sub>2</sub>SO<sub>4</sub>p, KOH, Mayer, Dragendorff and Lieberman-Burchard reagent.

### Tools

The tools used in this study included rotary evaporator, waterbath, soxhlet, petri dish, volumetric pipette, erlenmeyer flask, measuring cup, syringe, stirring rod, plethysmometer, analytical balance, porcelain dishes, oven, desiccators, wood clamps, tube racks, drip plates, spatulas, and spirits burners

### Preparation and extraction

Plant determination was conducted at the herbarium of School of Life Sciences, Bandung Institute of Technology. *A. altilis* leaves were cleaned, chopped and dried. Phytochemical screening was performed to determine the content of secondary metabolites, such as alkaloids, flavonoids, tannins, phenolics, triterpenoids, steroids, quinones, monoterpenes, sesquiterpenes, and saponins. Determination of ash content, water and ethanol soluble compounds were also performed. The leaves were extracted using soxhlet method with methanol solvent at 64.7 °C. Each filtrate was separated and evaporated using a rotary evaporator at 45 °C.

**Table 2. Phytochemical Screening of *A. Altilis* Leaves**

Metabolites	Attached Leaves	Detached Leaves
Alkaloid	-	-
Flavonoid	+	+
Tannin	+	+
Phenolic	+	+
Monoterpene, sesquiterpene	+	+
Steroid	+	+
Triterpenoides	-	-
Quinone	+	+
Saponin	-	-

**Table 3. Average of Edema Volume**

Group	V <sub>0</sub>	t <sub>1</sub>	t <sub>2</sub>	t <sub>3</sub>	t <sub>4</sub>	t <sub>5</sub>	t <sub>6</sub>
Negative control	0.035	0.051	0.061	0.061	0.063	0.065	0.065
Positive control	0.026	0.031	0.033	0.037	0.037	0.035	0.031
Attached 50 mg	0.028	0.035	0.04	0.043	0.043	0.043	0.043
Attached 100 mg	0.028	0.042	0.042	0.042	0.042	0.042	0.042
Detached 100 mg	0.025	0.032	0.033	0.033	0.035	0.035	0.032
Detached 50 mg	0.025	0.033	0.037	0.037	0.037	0.038	0.033

*Anti-inflammatory activity assay*

Experimental animals used in this study were Wistar strain male rats (180-250 g). Male rats were selected to avoid the influence of hormonal change in inflammatory responses. The animals were acclimatized for two weeks before the experiment. The animals were classified into 6 groups, each group consisting of 3 rats, i.e., negative control, positive control, and 4 treatment group with different doses. The negative control group was given 3 ml/200 g body weight 2% PGA suspension orally, while the positive control group was given 0.9 mg/ 200 g body weight diclofenac sodium orally. The experimental doses were 50mg/kg and 100mg/kg body weight, for both detached and attached leaves. Inflammation was induced by administering carrageenan on rats paw. The volume of edema was measured, before and after the treatment.

**Results and Discussion**

The results of plant determination showed that the plants used in this study was *A. altilis*

(Parkinson) Fosberg. The yield of extracts were 11.12% and 13.05% for attached and detached leaves, respectively. Methanol was selected as the solvent since it has better polarity compared to other solvents. The temperature used in the extraction process was based on the boiling point of the solvent. The percentage of ash contents and water soluble compounds of both types of leaves were relatively similar. However, the ethanol soluble compounds was higher in detached leaves (12.44%)<sub>6</sub> compared to attached leaves (8.92%). The results can be seen in the Table 1. Determination of ash content was conducted to determine the mineral and metal content from the initial process to the formation of extracts. Phytochemical screening showed that both types of leaves contained flavonoids, tannins, phenolics, monoterpenes, sesquiterpenes, steroids, and quinones (Table 2).

The anti-inflammatory activity assay showed that there was a decrease in the volume of edema in all groups, except negative control.

**Table 4. Percentage of the Inhibition of Inflammation**

Group	V <sub>0</sub> (%)	T <sub>1</sub> (%)	t <sub>2</sub> (%)	t <sub>3</sub> (%)	t <sub>4</sub> (%)	t <sub>5</sub> (%)	t <sub>6</sub> (%)
Negative control	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Positive control	0.00	68.75	73.08	57.69	60.71	70.00	83.33
Attached 50 mg	0.00	56.25	53.85	42.31	46.43	50.00	50.00
Attached 100 mg	0.00	12.50	46.15	46.15	50.00	53.33	53.33
Detached 100 mg	0.00	12.50	69.23	69.23	64.29	66.67	76.67
Detached 50 mg	0.00	50.00	53.85	53.85	57.14	56.67	73.33




Figure 1. Percentage of Inflammation Over Observed Time

The smaller the average volume of edema, the better the anti-inflammatory effect of this particular compound. The largest inhibition of inflammation was observed in positive control group with 83.33% reduction of edema. There were differences in anti-inflammatory activity between detached and attached leaves. The inhibition of edema in attached leaves were 50% and 53.33% for the concentration of 50 mg and 100 mg, respectively. The greater inhibition was observed in the detached leaves, with 73.33% and 76.67% inhibition, for the concentration of 50 mg and 100 mg, respectively. Among the treatment group, the extract of detached leaves 100 mg showed the best anti-inflammatory activity. Nevertheless, the percentage of the reduction was still higher than that of diclofenac sodium.

Differences in anti-inflammatory activity between detached and attached leaves might be due to differences in ethanol soluble compounds. Previous study showed that phenolic compounds might be responsible for anti-inflammatory activity of *A. altalis* plants. Further study is needed to confirm and elucidate the structure of the active anti-inflammatory compound in *A. altalis*.<sup>8-13</sup>

### Conclusion

*A. altalis* leaves extracts showed good anti-inflammatory properties and has the potential for development of anti-inflammatory drug.

### Acknowledgements

None declared.

### Funding

None.

### Conflict of Interest

None declared.

### References

1. Chen L, Deng H, Cui H, *et al*. Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget*. 2017;9(6):7204–7218.
2. Martino JV, Van Limbergen J, Cahill LE. The role of carrageenan and carboxymethylcellulose in the development of intestinal inflammation. *Frontiers in Pediatrics*. 2017;5:96.
3. Okin D, Medzhitov R. Evolution of inflammatory diseases. *Current Biology*. 2012;22(17):R733–R740.
4. Welz AN, Emberger-Klein A, Menrad K. Why people use herbal medicine: insights

- from a focus-group study in Germany. *BMC Complementary and Alternative Medicine*. 2018;18(1):92.
5. Yin SY, Wei WC, Jian FY, *et al*. Therapeutic applications of herbal medicines for cancer patients. *Evidence Based in Complementary and Alternative Medicine*. 2013;2013:302426.
  6. Sairam S, Urooj A. Safety evaluation of *Artocarpus altilis* as pharmaceutical agent in Wistar rats. *Journal of Toxicology*. 2014;2014:980404.
  7. Liu G, Kennedy R, Greenshields DL, *et al*. Detached and attached arabidopsis leaf assays reveal distinctive defense responses against hemibiotrophic *Colletotrichum spp.* *Molecular Plant Interaction*. 2007;20:1308–1319.
  8. Nwokocha CR, Owu DU, McLaren M, *et al*. Possible mechanisms of action of the aqueous extract of *Artocarpus altilis* (breadfruit) leaves in producing hypotension in normotensive sprague-dawley rats. *Pharmaceutical Biology*. 2012;50(9):1096-1102.
  9. Riasari H, Sukrasno S, Ruslan H. Metabolite profile of various development bread fruit leaves (*Artocarpus altilis*. Parkinson. Fosberg) and the identification of their major componens. *International Journal of Pharmaceutical Sciences and Research*. 2015;6(5):2170-2177.
  10. Turi CE, Liu Y, Ragone D. Breadfruit (*Artocarpus altilis* and hybrids): A traditional crop with the potential to prevent hunger and mitigate diabetes in Oceania. *Trends in Food Science and Technology*. 2015;45:264–272.
  11. Yao X, Wu D, Dong N, *et al*. Moracin C, A Phenolic compound isolated from *Artocarpus heterophyllus*, suppresses lipopolysaccharide-activated inflammatory responses in murine. *International Journal of Molecular Science*. 2016;17(8):1199.
  12. Lan WC, Tzeng CW, Lin CC. Prenylated flavonoids from *Artocarpus altilis*: Antioxidant activities and inhibitory effects on melanin production. *Phytochemistry*. 2013;89:78–88.
  13. Tiraravesit N, Yakaew S, Rukchay R, *et al*. *Artocarpus altilis* heartwood extract protects skin against UVB in vitro and in vivo. *Journal of Ethnopharmacology*. 2015;175:153–162.

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

%

INTERNET SOURCES

19%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Nur Azizah, Eli Halimah, Irma Melyani Puspitasari, Aliya Nur Hasanah. "Simultaneous Use of Herbal Medicines and Antihypertensive Drugs Among Hypertensive Patients in the Community: A Review", Journal of Multidisciplinary Healthcare, 2021  
Publication 2%
- 2 M. B. A. Kadir, E. R. Kasimo. "The Effect of Breadfruit Leaf Extract (Artocarpus Altilis P) on Spatial Memory of in Male Strain Mice which Indicated by Alcohol", Journal for Quality in Public Health, 2020  
Publication 2%
- 3 Lee-Hoon Ho, Shi-Yun Wong. "Resistant Starch from Exotic Fruit and Its Functional Properties: A Review of Recent Research", IntechOpen, 2019  
Publication 2%
- 4 Bai-Luh Wei, Jing-Ru Weng, Pao-Hui Chiu, Chi-Feng Hung, Jih-Pyang Wang, Chun-Nan Lin. "Antiinflammatory Flavonoids from and ", 1%

5

J. Debode. "Quantitative detection and monitoring of *Colletotrichum acutatum* in strawberry leaves using real-time PCR", *Plant Pathology*, 06/2009

Publication

---

1 %

6

Kyung-Mi Lee, Taejin Park, Min-Seon Kim, Jin-Soo Park, Won-Jae Chi, Seung-Young Kim. "Anti-inflammatory Activities of 7,8-Dihydroxy-4-Methylcoumarin Acetylation Products NF- $\kappa$ B and MAPK Pathways in LPS-Stimulated RAW 264.7 Cells ", *Natural Product Communications*, 2022

Publication

---

1 %

7

Wen-Jing Yang, Ji-Xiang He, Ming-Xing Zhou, Min Huang, Shu-Qi Wang, Xiao-Ning Wang, Hong-Xiang Lou, Dong-Mei Ren, Tao Shen. "An isopentenyl-substituted flavonoid norartocarpin activates Nrf2 signalling pathway and prevents oxidative insults in human lung epithelial cells", *Free Radical Research*, 2019

Publication

---

1 %

8

Yuhui Ye, Dongxiao Sun-Waterhouse, Lijun You, Arshad Mehmood Abbasi. "Harnessing food-based bioactive compounds to reduce

1 %



the effects of ultraviolet radiation: a review exploring the link between food and human health", International Journal of Food Science & Technology, 2017

Publication

---

9

Nokulunga Hlengwa, Christo J. F. Muller, Albertus K. Basson, Sandra Bowles, Johan Louw, Charles Awortwe. "Herbal supplements interactions with oral oestrogen - based contraceptive metabolism and transport", Phytotherapy Research, 2020

Publication

---

1 %

10

Maged Younes, Peter Aggett, Fernando Aguilar, Riccardo Crebelli et al. "Re - evaluation of carrageenan (E407) and processed Eucheuma seaweed (E407a) as food additives", EFSA Journal, 2018

Publication

---

1 %

11

Merna G. Aboismaiel, Mohamed El-Mesery, Amro El-Karef, Mamdouh M. El-Shishtawy. "Hesperetin upregulates Fas/FasL expression and potentiates the antitumor effect of 5-fluorouracil in rat model of hepatocellular carcinoma", Egyptian Journal of Basic and Applied Sciences, 2020

Publication

---

1 %

12

Ina Vandebroek, David Picking. "Popular Medicinal Plants in Portland and Kingston,

1 %

13

N Nurmalasari, Y Yulizar, D O B Apriandanu. "Bi O nanoparticles: synthesis, characterizations, and photocatalytic activity", IOP Conference Series: Materials Science and Engineering, 2020

Publication

---

1 %

14

Price, Jordan V., David J. Haddon, Dodge Kemmer, Guillaume Delepine, Gil Mandelbaum, Justin A. Jarrell, Rohit Gupta, Imelda Balboni, Eliza F. Chakravarty, Jeremy Sokolove, Anthony K. Shum, Mark S. Anderson, Mickie H. Cheng, William H. Robinson, Sarah K. Browne, Steven M. Holland, Emily C. Baechler, and Paul J. Utz. "Protein microarray analysis reveals BAFF-binding autoantibodies in systemic lupus erythematosus", Journal of Clinical Investigation, 2013.

Publication

---

1 %

15

"Ethnobotany of the Mountain Regions of Southeast Asia", Springer Science and Business Media LLC, 2021

Publication

---

<1 %

16

Cheilane T. de Souza, Sarah A.R. Soares, Antonio F.S. Queiroz, Ana M.P. dos Santos,

<1 %

Sergio L.C. Ferreira. "Determination and evaluation of the mineral composition of breadfruit (Artocarpus altilis) using multivariate analysis technique", *Microchemical Journal*, 2016

Publication

---

17

Amanda de Assis Carneiro, Simone Batista Pires Sinoti, Marcela Medeiros de Freitas, Luiz Alberto Simeoni et al. "Hydroethanolic Extract of *Morusnigra* L. Leaves: A Dual PPAR- $\alpha/\gamma$  Agonist with Anti-Inflammatory Properties in Lipopolysaccharide-Stimulated RAW 264.7", *Plants*, 2022

Publication

---

<1 %

18

Arunaksharan Narayanankutty. "Pharmacological Potentials and Nutritional Values of Tropical and Subtropical Fruits of India: Emphasis on their Anticancer Bioactive Components", *Recent Patents on Anti-Cancer Drug Discovery*, 2022

Publication

---

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

# Hesti Riasari\_PCPR

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---