

BERKAS JURNAL : 2022

Link jurnal : <https://www.phcogj.com/article/1396>

Pharmacological Activities and Phytochemical Compounds: Overview of Pouteria Genus

Halaman Sampul Jurnal Online :

The screenshot shows the homepage of the Open Access Macedonian Journal of Medical Sciences. The header is dark red with white text for navigation: "Open Access Macedonian Journal of Medical Sciences", "Journal Info", "Announcements", "Policies", "For Authors", "For Reviewers", "Issues", "Categories", "Membership", and a search icon. Below the header, there's a breadcrumb "Home / About the Journal" and a main heading "About the Journal". A cover image of the journal is displayed, featuring the "oa mjms" logo and the ISSN 1857-0655. The main text describes the journal as a top-tier open access medical science journal published by the Scientific Foundation SPIROSKI, covering various areas of medical science. On the right side, there are sections for "Submit in our journals" listing several journals like SEEJAD, SEEJCA, SEEJIM, SEEJHG, and MMEJ, and a "Co-publishing" section with a Sciendo logo and a woman's portrait. At the bottom right, a "Scientific Publications" section highlights Scopus and Web of Science journals with a Hirsch Index increase.

Open Access Macedonian Journal of Medical Sciences

Journal Info ▾ Announcements ▾ Policies ▾ For Authors ▾ For Reviewers ▾ Issues ▾ Categories ▾ Membership ▾ [Search](#)

[Home](#) / [About the Journal](#)

About the Journal



Open Access Macedonian Journal of Medical Sciences (OAMJMS) [formerly known as Macedonian Journal of Medical Sciences] is a top-tier open access medical science journal published by the Scientific Foundation SPIROSKI, Rajko Zhinzifov No 48, 1000 Skopje, Republic of Macedonia. *OAMJMS* is an international, modern, general medical journal covering all areas in the medical sciences, from basic studies to large clinical trials and cost-effectiveness analyses. We publish mostly human studies that substantially enhance our understanding of disease epidemiology, etiology, and physiology; the development of prognostic and diagnostic technologies; trials that test the efficacy of specific interventions and those that compare different treatments; dental sciences; nursing; and systematic reviews. We aim to promote translation of basic research into clinical investigation, and of clinical evidence into practice. We publish occasional studies in animal models when they report outstanding research findings that are highly clinically relevant. Our audience is the international medical community as well as educators, policy makers, patient advocacy groups, and interested members of the public around the world. *OAMJMS* is published 4 times per year.

Submit in our journals

- [South East European Journal of Architecture and Design \(SEEJAD\)](#)
- [South East European Journal of Cardiology \(SEEJCA\)](#)
- [South East European Journal of Immunology \(SEEJIM\)](#)
- [South East European Journal of Human Genetics \(SEEJHG\)](#)
- [Macedonian Medical Electronic Journal \(MMEJ\)](#)

Co-publishing

 sciendo
Your publishing needs met. 

Scientific Publications


Publication in **Scopus** and **Web of Science** Journals
+ Hirsch Index Increase

Dewan Redaksi/Editorial Board :

Open Access Macedonian Journal of Medical Sciences

[Journal Info](#) - [Announcements](#) - [Policies](#) - [For Authors](#) - [For Reviewers](#) - [Issues](#) - [Categories](#) - [Members](#)

[Home](#) / [Editorial Team](#)

Editorial Team

Journal Manager

Eli Djulejic, MD, Open Access Macedonian Journal of Medical Sciences, Belgrade, Serbia

elidjulejic@yahoo.com

ORCID ID: <https://orcid.org/0000-0002-1736-9029>

Web of Science ResearcherID: AAC-3225-2020

Scopus Author ID: 33767574400

Editor-in-Chief

Mirko Zhivko Spiroski, MD, PhD, Scientific Foundation SPIROSKI, Rajko Zhinzifov No 48, 1000 Skopje, Republic of Macedonia

mirkospiroski@foundationspiroski.eu

ORCID ID: <https://orcid.org/0000-0001-5559-7981>

Web of Science ResearcherID: F-7606-2012

Scopus Author ID: 6602348069

Section Editors (Deputy Editors-in-Chief)

Stoleski Sasho, MD, PhD, Institute for Occupational Health of Republic of Macedonia - Skopje, WHO Collaborating Center, GAZLEN Collaborating Center, II Makedonska brigada 43, 1000 Skopje, Republic of Macedonia

sstoleski@yahoo.com

ORCID ID: <https://orcid.org/0000-0003-1278-903X>

Web of Science ResearcherID: B-2304-2015

Scopus Author ID: 9943962300

Slavica Hristomanova-Mitkovska, MD, MSc, Institute of Human Genetics, University Medical Center Göttingen, Heinrich-Düker-Weg 12, 37079 Göttingen, Germany

sacka_h@yahoo.com

ORCID ID: <https://orcid.org/0000-0002-7770-5055>

Web of Science ResearcherID: G-8801-2012

Scopus Author ID: 33767823700

Igor Spiroski, MD, University Clinic of Cardiology, Faculty of Medicine, Ss Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia

ispiroski@gmail.com

ORCID ID: <https://orcid.org/0000-0002-3448-0685>

Web of Science ResearcherID: AAA-5652-2020

Scopus Author ID: 23971606200

Ksenija Bogoeva-Kostovska, MD, PhD, PHO Prof Bogoev, Skopje, Republic of Macedonia

ksenijabogoeva@gmail.com

Sinisha Stojanovski, MD, PhD. Medical Faculty, Ss Cyril and Methodius University of Skopje,
Skopje, Republic of Macedonia
sinisa.stojanovski@hotmail.com
ORCID ID: <https://orcid.org/0000-0002-4967-2102>
Web of Science ResearcherID: [AFL-9514-2022](https://orcid.org/0000-0002-4967-2102)
Scopus Author ID: [6504203064](https://orcid.org/0000-0002-4967-2102)

Filip Koneski, Doctor of Dental Medicine, MSc. University Clinic for Maxillofacial Surgery in Skopje, Ss. Cyril and Methodius
University of Skopje, Faculty of Dental Medicine, Skopje, Republic of Macedonia
dr.koneski@gmail.com
ORCID ID: <https://orcid.org/0000-0003-2412-7594>
Web of Science ResearcherID: [J-9194-2019](https://orcid.org/0000-0003-2412-7594)
Scopus Author ID: [57073856400](https://orcid.org/0000-0003-2412-7594)

Aleksandar Iliev, Doctor of Dental Medicine, PhD. Department of Maxillofacial Surgery, Faculty of Dentistry, Ss Cyril and
Methodius University of Skopje, Skopje, Republic of Macedonia
aleksandar.iliev@gmail.com
ORCID ID: <https://orcid.org/0000-0001-5348-9143>
Web of Science ResearcherID: [ACV-7553-2022](https://orcid.org/0000-0001-5348-9143)
Scopus Author ID: [57021392000](https://orcid.org/0000-0001-5348-9143)

Layout Editor and Electronic Publishing

Ivo Spiroski, Eng, MSc. Scientific Foundation SPIROSKI, Rajko Zhinzifov No 48, 1000 Skopje, Republic of Macedonia
ivospiroski@foundationspiroski.eu
ORCID ID: <https://orcid.org/0000-0003-4518-2138>
Web of Science ResearcherID: [F-2799-2014](https://orcid.org/0000-0003-4518-2138)

Production Editor

Teodora Fildishevska, Scientific Foundation SPIROSKI, Skopje, Republic of North Macedonia
E-mail: teodora.fildishevska@sfsmdr.mk
ORCID ID: <https://orcid.org/0000-0003-1483-4789>

Copyeditor

Sinjore, India

Editorial Board

Nikola Angelov, DDS, MS, PhD. Professor and Chair, UTHealth School of Dentistry, Loma Linda University School of
Dentistry, Department of Periodontics, Loma Linda, CA, 92350, United States
nangelov@llu.edu
ORCID ID: <https://orcid.org/0000-0002-0800-2778>

Ramush Bejtiqi, MD, PhD. University Clinical Centre of Kosovo, Paediatric Clinic, Kosovo
rbejtiqi@hotmail.com
ORCID ID: <https://orcid.org/0000-0002-5580-4458>

Semra Čavaljuga, MD, PhD. Department of Epidemiology and Biostatistics, Faculty of Medicine, Sarajevo, Bosnia and
Herzegovina
scavaljuga@gmail.com

Editorial Board

Nikola Angelov, DDS, MS, PhD. Professor and Chair, UTHealth School of Dentistry, Loma Linda University School of Dentistry, Department of Periodontics, Loma Linda, CA, 92350, United States

nangelov@llu.edu

ORCID ID: <https://orcid.org/0000-0002-0800-2778>

Ramush Bejqi, MD, PhD. University Clinical Centre of Kosovo, Paediatric Clinic, Kosovo

rbejqi@hotmail.com

ORCID ID: <https://orcid.org/0000-0002-5580-4458>

Semra Čavaljuga, MD, PhD. Department of Epidemiology and Biostatistics, Faculty of Medicine, Sarajevo, Bosnia and Herzegovina

scavaljuga@gmail.com

ORCID ID: <https://orcid.org/0000-0003-0370-9560>

Pei-Yi Chu, MD. Diagnostic and research pathologist, Department of Surgical Pathology, Changhua Christian Hospital, Taiwan. Address: 135 Nan-Shiao Street, Changhua 500-06, Taiwan, Province of China

chupeiyl@yahoo.com.tw

ORCID ID: <https://orcid.org/0000-0001-7336-2465>

Ivo Donkov, MD, PhD. Staff Urologist, Lincoln County Hospital, United Kingdom

idonkov@hotmail.com

ORCID ID: <https://orcid.org/0009-0007-5669-8734>

Andrew J. Dwork, MD, PhD. Departments of Pathology and Cell Biology and Psychiatry, College of Physicians and Surgeons of Columbia University; Division of Molecular Imaging and Neuropathology, New York State Psychiatric Institute, Unit 62, 722 West 168th Street, New York, NY 10032, United States

ajd6@columbia.edu

ORCID ID: <https://orcid.org/0000-0002-9862-0384>

Adriana Galan, Department of Health Programmes and Health Promotion, Institute of Public Health, Bucharest, Romania

agalan@ispb.ro

Tania Santos Giani, MD, PhD. Estacio de Sa University, in Health Sciences, Brazil

tianasantosgiani@gmail.com

ORCID ID: <https://orcid.org/0000-0002-1913-5090>

Iva Ivanovska, MD, PhD. Harvard Medical School, Department of Genetics, 77 Avenue Louis Pasteur, NRB room 239, Boston, MA 02115, United States

ivanovska@genetics.med.harvard.edu

ORCID ID: <https://orcid.org/0000-0002-0633-5333>

Jerzy Jablęcki, MD, PhD. Division of General Surgery St. Jadwiga of Silesia Hospital, Trzebnica; Head, Subdepartment of Hand Surgery and Replantation St. Jadwiga of Silesia Hospital, Trzebnica; Professor, Department of Public Health, State Higher Professional Medical School, Opole, Poland. 55-100 Trzebnica, ul. Prusicka 53, Poland

jerzy.jablęcki@interia.pl

ORCID ID: <https://orcid.org/0000-0002-9498-1051>

Mehrdad Jalalian Hosseini, MD, PhD. Khorasan-e Razavi Blood Center, Mashhad, Iran, Islamic Republic of

Mehrdad.medic@gmail.com

Radka Kaneva, PhD. Department of Medical Chemistry and Biochemistry, Medical University - Sofia, Bulgaria

kaneva64@yahoo.com

ORCID ID: <https://orcid.org/0000-0002-1437-1393>

Kostandina Leonida Korneti-Pekavska, MD, PhD. Ss Cyril and Methodius University of Skopje, Faculty of Medicine, Skopje, Republic of Macedonia

kostandina_korneti@yahoo.com

DAFTAR ISI :

Home / Archives / Vol. 10 No. A (2022): A - Basic Sciences

Vol. 10 No. A (2022): A - Basic Sciences



Published: 2022-01-01

Genetics

Role of MicroRNAs in the Development of Chronic Liver Disease in Hepatitis Virus-Infected Egyptian Population

DOI: <https://doi.org/10.3889/oamjms.2022.10902>

Marwa Hassan, Eman El-Ahwany, Mohamed Elzallat, Ali Abdel Rahim, Hoda Abu-Taleb, Yosry Abdelrahman, Moataz Hassanein (Author) 1555-1559



Relationship Between Phylogenetic of Apium and Foeniculum Plants from Central Java, Indonesia, and Their Secondary Metabolites Potency against COVID-19 Protease

DOI: <https://doi.org/10.3889/oamjms.2022.9852>

Hermin Pancasakti Kusumaningrum, Rejeki Siti Ferniah, Siti Nur Jannah, Mufida Budi Kurniawati, Anis Afifah, Yoshua Mario Sumbodo, Salma Seftia Hanif, Tia Erfianti, Yuriza Eshananda (Author) 1234-1241



Genetic Study of Chemokine Ligand 1 in Colorectal Carcinoma using Quantitative Real-Time PCR

DOI: <https://doi.org/10.3889/oamjms.2022.8645>

Athraa Alshimerry, Dalia Amer Khudhair, Roaa Salih Mahdi (Author) 656-660



Low-dose Indonesian Aloe vera Increases Viability and Migration of the Fibroblast: An In Vitro Study

DOI: <https://doi.org/10.3889/oamjms.2022.8177>

Januar Rizqi, Akbar Satria Fitriawan (Author)

256-261



Determination of Total Phenolic Content, Analysis of Bioactive Compound Components, and Antioxidant Activity of Ethyl Acetate Seri (Muntingia calabura L.) Leaves from North Sumatera Province, Indonesia

DOI: <https://doi.org/10.3889/oamjms.2022.8362>

Risanti Febrine Ropita Situmorang, Kasta Gurning, Vera Estefania Kaban, Mastiur Julianti Butar-Butar, Susi Amenta Beru Perangin-Angin (Author)

240-244



Cost-Effectiveness of Ticagrelor for Acute Coronary Disease to Prevent Cardiovascular Events in Three Hospitals in Indonesia

DOI: <https://doi.org/10.3889/oamjms.2022.9671>

Jarir At Thobari, Lucia Krisdinarti, Dhite Nugroho, Jonathan Haposan, Isman Firdaus, Rr. Arum Ramadhyan Suryandani, Muhammad Munawar, Jimmy Agung (Author)

1211-1217



Effect of Different Solvent on Phytochemical Content, Tyrosinase Inhibition and Antioxidant Activities of Campolay (*Pouteria campechiana* kunth. [Baehni.]

DOI: <https://doi.org/10.3889/oamjms.2022.8204>

Sani Nurlaela Fitriansyah, Rika Hartati, Irda Fidrianny (Author)

158-163



Antidiabetic Effect Test of Insulin Stem Extract (*Tithonia diversifolia*) Toward Streptozotocin-Induced Diabetic Rats (*Rattus Norvegicus*)

DOI: <https://doi.org/10.3889/oamjms.2022.8411>

Suherman Suherman, Baharuddin Hamzah, Sri Hastuti V. Pulukadang, Sitti Rahmawati, Muhammad Fakhrol Hardani, Ririen Hardani, Andi Saifah (Author)

1006-1010



Unused Parts of Jackfruit (*Artocarpus heterophyllus*): Prospective In Vitro Antioxidative Activity



Effect of Different Solvent on Phytochemical Content, Tyrosinase Inhibition and Antioxidant Activities of Campolay (*Pouteria campechiana* kunth. [Baehni.]

Sani Nurlaela Fitrianyah^{1,2*} , R. Hartati¹, I. Fidrianny¹

¹Department of Pharmaceutical Biology, School of Pharmacy, Bandung Institute of Technology, Bandung, Indonesia; ²Department of Pharmaceutical Biology, Indonesian School of Pharmacy (Sekolah Tinggi Farmasi Indonesia), Bandung, Indonesia

Abstract

BACKGROUND: *Pouteria campechiana* leaves are reported to have phenol and flavonoid compounds. Phenolic and flavonoid compounds can act as tyrosinase inhibitor and antioxidant.

AIM: The purpose of this study was to compare phytochemical content, tyrosinase inhibition, antioxidant activities, and determine of marker compound from *P. campechiana* leaves extract with different polarities solvent.

METHODS: In addition, the content of marker compound from *P. campechiana* leaves extract was determined by HPLC.

RESULTS: The highest total phenolic content (TPC) 7.83 GAE/100 g extract, IC₅₀ of tyrosinase 171.512 ± 1.352 and IC₅₀ of DPPH 0.968 ± 0.008 was given by ethanolic extract (DE). Meanwhile ethyl acetate extract (DEA) had the highest of total flavonoid content 2.544 ± 0.554 QE/100 g extract. The TPC correlated with tyrosinase inhibitory activity and antioxidant activity.

CONCLUSION: Quercitrin was as marker compound from *P. campechiana* leaves extract, and quercitrin content in the DEA of *P. campechiana* leaves was 3.539%, while in the DE was 0.153%.

Edited by: Sinisa Stojanoski
Citation: Fitrianyah SN, Hartati R, Fidrianny I. Effect of Different Solvent on Phytochemical Content, Tyrosinase Inhibition and Antioxidant Activities of Campolay (*Pouteria campechiana* kunth. [Baehni.]). Open Access Maced J Med Sci. 2022 Jan 18; 10(A):158-163. https://doi.org/10.3889/oamjms.2022.8204
Keywords: *Pouteria campechiana*; Tyrosinase; Antioxidant; Quercitrin
***Correspondence:** Sani Nurlaela Fitrianyah, Department of Pharmaceutical Biology, School of Pharmacy, Bandung Institute of Technology, Bandung, Indonesia. E-mail: saninurlaela@stfi.ac.id
Received: 12-Sep-2021
Revised: 29-Dec-2021
Accepted: 08-Jan-2022
Copyright: © 2022 Sani Nurlaela Fitrianyah, R. Hartati, I. Fidrianny
Funding: This study was supported by the Indonesia Endowment Fund for Education
Competing Interests: The authors have declared that no competing interests exist
Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Introduction

Exposure of UV light can produce Reactive Oxidative Singlet (ROS). If the mechanism action of ROS is not inhibited, it will increase the tyrosinase action [1]. Tyrosinase is enzyme that plays the role in biosynthesis of melanin in plant and mammalian cell, including human skin [2]. The mechanism action of tyrosinase enzyme divided into two, namely monophenolase activity using L-tyrosine substrate and diphenolase using L-DOPA substrate [2], [3]. Diphenolase activity will produce ortho-quinone, meanwhile melanin will be produced by non-enzymatic reaction. This process is called melanogenesis [4].

Antioxidants compounds can inhibit the reaction of free radicals. An example of a free radical is ROS which is found in the skin. When antioxidant can inhibit reaction of ROS which is caused by UV light exposure on the skin, it will reduce tyrosinase enzyme activity. Phenolic acid [5], flavonol [6], and stilbenoide [7] were antioxidant compounds as well as tyrosinase inhibitors.

Medicinal plants are a source of raw material which contain phenolic, flavonoids and other polyphenols. The biological activity of a plant can be caused by the presence of phytochemical compound. The solvent used and extraction method will affect the phytochemical compound resulting. *Pouteria campechiana* is one of the medicinal plants in Sapotaceae family. In Indonesia, *P. campechiana* was called as campolay. *P. campechiana* fruits extract contains gallic acid, myricitrin and catechin which act as antioxidant compounds [8]. In addition, taxifolin, quercetin glycoside and myricetin glycoside were found in *P. campechiana* leaves which its toxicity had been tested [9]. The comparison of chemical content, tyrosinase inhibitory activity, antioxidant activity of *P. campechiana* leaves extract have not been reported with different solvent polarity levels.

In this study, the total chemical content was calculated, inhibitory of tyrosinase and antioxidant activities were tested, then marker compound from selected extract of *P. campechiana* was identified and determined.

Experimental

Material

P. campechiana leaves were collected from Bandung, West Java, Indonesia in December 2020.

Chemical material

DPPH (Sigma Aldrich), tyrosinase enzyme (T3824-25KU, Sigma Aldrich), L-DOPA (Sigma Aldrich), quercetin, kaemferol, apigenin, gallic acid, chlorogenic acid, luteolin 7-O glycoside, and rutin which were obtained from Sigma Aldrich, quercitrin (Markherb), methanol pro analysis, ethanol 96%, ethyl acetate, n-hexane, Folin–Ciocalteu reagent, aluminum chloride and other chemical materials which used in this study were analytical grade.

Preparation extract

Extraction was conducted using Soxhlet with increasing polarity solvent, ranging from nonpolar (n-hexane), semi-polar (ethyl acetate), and polar solvent (ethanol 96%). 400 g of *P. campechiana* leaves powder was extracted with n-hexane, the filtrate and residue are separated. The residue was dried, and then extracted with ethyl acetate. Finally, the residue was extracted using ethanol 96%. Each extract was thickened using rotary evaporator with a temperature of 50°C to produce n-hexane extract (DN), ethyl acetate extract (DEA), and ethanol 96% extract (DE). Each extraction process with Soxhlet was carried out triplicate. The use of this extraction method is expected to be more effective in separating the phytochemical content.

Total phenolic content (TPC)

Determination of TPC using modification of Pourmorad's method [10]. Each extract was tested triplicate. The test was measured at a wavelength of 765 nm. TPC was calculated by gallic acid calibration curve with concentration in the range of 10-250 µg/ml and expressed as gallic acid equivalent (GAE) per 100 g extract (g GAE/100 g extract).

Total flavonoid content (TFC)

Determination of TFC using of Chang's method modified [11]. Each extract was tested triplicate. The test was investigated at a wavelength of 415 nm. TFC was calculated by quercetin calibration curve with concentration ranged from 25 to 250 µg/ml and presented as quercetin equivalent (QE) per 100 g extract (g QE/100 g extract).

Tyrosinase inhibitory activity

Determination of tyrosinase inhibitory activity using as spectrophotometer in 96-well plates (Biologix®) following the procedure of Masuda's method [12] with slight modification. Tyrosinase inhibitory activity was determined by diphenolase mechanism, with L-Dopa as substrate. Standard of tyrosinase inhibitor (kojic acid) were dissolved in phosphate buffer (pH 6.8) which contained DMSO 5%. The enzyme concentration used was 300 unit/ml. Absorbance was investigated at λ 475 nm. Concentration of DN, DEA and DE 1000 µg/ml were carried out for testing tyrosinase inhibitory activity. Then, 10 concentration series was prepared from the selected sample to find the IC₅₀ value.

Antioxidant activity

Determination of antioxidant activity was used modified Blois's method [13]. The free radical DPPH with a concentration of 39.4 µg/ml in methanol. The standard was ascorbic acid. Absorbance measurement at λ 517 nm using a UV-Vis spectrophotometer (Beckman Coulter DU 720). Extracts and standard were performed in three replications. The final parameter used as antioxidant activity was antioxidant activity index (AAI), which applied the formula $AAI = (\text{Final concentration of DPPH } [\mu\text{g/ml}]/IC_{50} (\mu\text{g/ml}))$ [14].

Statistical analysis

The results were presented with the mean ± SD value from three repetitions of the test, using MS excel software to evaluate the results of the IC₅₀ value. Statistical analysis was also used to see the correlation between the TPC and TFC to tyrosinase inhibitory and antioxidant activities with Pearson's method in Minitab-19 application.

Identification of marker compound from selected extract

Identification of marker compound was carried out qualitatively. The first method using TLC co-chromatography to compare the Retardation factor (Rf) values with various standard compounds. The second method used was HPLC (LC-20AD) by comparing with quercitrin as standard compound selected based on the TLC co-chromatography. The stationary phase in HPLC used LiChrospher® 100 RP-C18 5 µm (length 100 mm, diameter 4 mm, 20 mm in column [Merck]). The mobile phase used was 0.01% H₃PO₄ (eluent A) dan methanol (eluent B). Eluent B, 2.5 min 40%, 2.51 to 7 min 60%, 7.01-12 min 70%, and 40% for linier gradient from 12 to 15 min. Flow rate used 1 ml/min, (CTO-20A pump, Shimadzu, Japan). Injection volume 20 µl. Detector used 366 nm (Detector UV/Vis SPD-20A, Shimadzu, Japan). The parameter used was

retention time from standard compound (quercitrin) and sample. Sample and quercitrin were diluted in methanol with 50.000 µg/ml for DE, 10.000 µg/ml for DEA and 500 µg/ml for quercitrin.

Determination of marker content

Determination of marker content from selected extract used the one-point method. Calculation the content of marker was by comparing the area under curve (AUC) of marker in extract and AUC of standard compound.

Results and Discussion

Extraction of *P. campechiana* leaves

Extraction was carried out by Soxhlet with different polarity solvents. The compound in the crude drug will be derived based on polarity. The yield obtained from DN, DEA and DEs of *P. campechiana* leaves was showed in Table 1. The yield of the extracts indicated amount of the secondary metabolite which was extracted in each solvent with different polarity. DE had the largest yield compared to DN and DEA. This result showed that extraction of *P. campechiana* leaves by Soxhlet with ethanol 96% could attract more metabolite from *P. campechiana* leaves.

Table 1: The yield of *Pouteria campechiana* leaves extracts

Sample	Yield of extract (g extract/100 g dried leaves)
DN	7.24
DEA	5.25
DE	23.00

DN: n-hexane extract, DEA: Ethyl acetate extract, DE: Ethanol extract.

Determination of phytochemical content

Determination of phytochemical content was conducted for TPC and TFC. This is based on phenolic and flavonoid plays a role in tyrosinase inhibitory and antioxidant activities.

TPC was calculated using gallic acid curve with linier regression equation $y = 0.0062x + 0.0592$; $R^2 = 0.999$ (Figure 1) and presented by g GAE/100 g extract. It has no report regarding TPC and TFC in *P. campechiana* leaves extract using the multilevel extraction method by Soxhlet. DE had higher TPC and significantly different compared to DN and DEA ($p < 0.05$). The TPC of ethanol leaves extract was higher than DN and DEAs (Figure 3). Phenolic compound is a group of secondary metabolites in plants. The phenolic group included polyphenols. Polyphenols included flavonoids, phenolic acids, stilbenoid and lignan [15]. The phenolic group which has more OH groups showed high polarity and will be more effective if extracted with a




Figure 1: Calibration curve of gallic acid

polar solvent, such as ethanol. The fewer OH groups in phenolic group will be extracted with semipolar solvents such as ethyl acetate.

The TFC was determined based on the quercetin standard curve [Figure 2], $y = 0.0061x + 0.0796$; $R^2=0.994$ and exposed with g QE/100 g extract. DEA showed higher TFC and significantly different compared to DN and DE ($p < 0.05$). It showed the DEA had the highest TFC compared to DE and DNs. This means, that ethyl acetate solvent was more effective for extracting flavonoids from *P. campechiana* leaves. Based on the structure, the flavonoid group can be divided into flavones group such as apigenin, flavonols group such as quercetin, quercitrin and myricitrin, flavan group, flavanol and anthocyanidin group [16].

Tyrosinase inhibitory activity

The tyrosinase inhibitory activity of *P. campechiana* leaves extract was expressed in Table 2. Determination of tyrosinase inhibitory activity using modification method of Masuda [12]. Kojic acid was used as tyrosinase inhibitor. Inhibitory of tyrosinase activity from DE, DEA and DNs of *P. campechiana* leaves were the first time reported. The result of the study showed that DE had the highest tyrosinase inhibitory activity compared to DEA and DN. The DN had lowest of tyrosinase inhibitory activity which was 15.078 ± 1.785 at 1000 µg/ml. Therefore, DN was not continued to determine the IC_{50} tyrosinase inhibitory value. The difference in tyrosinase inhibitory activity in DE, DEA and DN extract could be due to differences in phytochemical content that extracted in each extract. Extraction methods with different solvent polarity levels can extract different chemical constituents. Differences in the chemical structure of each compound will affect the strength of tyrosinase inhibitory and antioxidant activities [17]. Phenols and flavonoids can contribute to antioxidant and tyrosinase inhibitory activity [18]. The number of hydroxy groups can increase antioxidant activity and tyrosinase inhibitory activity [1]. According to the literature, Manilkara zapota belong to Sapotaceae family gave tyrosinase inhibitory activity. The chemical constituent of Manilkara zapota belonging to the triterpenoids, sterols, flavonoids, and phenolic groups presented tyrosinase inhibitory activity with

Table 2: IC₅₀ of DPPH and IC₅₀ of Tyrosinase *Pouteria campechiana* leaves extract

Extract	IC ₅₀ DPPH (µg/ml)	IC ₅₀ Tyrosinase (µg/ml)
DN	8.602 ± 1.226	-
DEA	1.040 ± 0.101	828.537 ± 0.962
DE	0.968 ± 0.008	176.553 ± 4.613
Ascorbic acid	0.570 ± 0.002	-
Kojic acid	-	10.513 ± 0.707

DN: n-hexane extract, DEA: Ethyl acetate extract, DE: Ethanol extract.





Figure 2: Calibration curve of quercetin

Figure 3: Phytochemical content of *Pouteria campechiana* leaves extract

monophenolase and diphenolase. Triterpenoids showed strong inhibitory of tyrosinase activity with monophenolase mechanism. Myricitrin which was a flavonoids glycoside from *Manilkara zapota* leaves extract, showed weak of tyrosinase inhibitory activity [19].


Antioxidant activity

Antioxidant activity of *P. campechiana* leaves extract can be seen in Table 3 and Figure 4. The value of antioxidant activity was presented in IC₅₀ and AAI. Ascorbic acid was used as standard. DPPH used as free radical. According to Scherer and Godoy [14], if AAI < 0.5 was weak antioxidant, 0.5–1 medium antioxidant, 1–2 strong antioxidant and more than 2 classified as very strong antioxidant. The highest AAI value was

Table 3: Correlation of phytochemical content with tyrosinase inhibitory and antioxidant activities

Activity parameter	Pearson's correlation coefficient (r)	
	TPC	TFC
AAI	0.981**	-0.978**
IC ₅₀ tyrosinase	-1.00**	0.999**

TPC: Total phenolic content, TFC: Total flavonoid content, AAI: Antioxidant activity index.

Figure 4: AAI of *Pouteria campechiana* leaves extract

in DE, followed by DEA and finally by DN extract of *P. campechiana* leaves. However, if classification based on the AAI value, the DE, DEA and DN of *P. campechiana* leaves were classified as very strong antioxidant. The AAI of DE, DEA, and DN was lower than the AAI of ascorbic acid. The phenolic group can contribute to antioxidant activity [20]. The position of the OH group on flavonoids can affect the antioxidant activity. The presence of OH groups in ring B, at C-3 and ortho-hydroxy groups in ring A and B can increase antioxidant activity. In addition to the OH group, the presence of a double bond at the C2-C3 position, and the presence of a carbonyl group at C4 can also increase antioxidant activity [16].

Correlation of total phenolic and flavonoid content with tyrosinase inhibitory and antioxidant activities

TPC had a positive and significant correlation with AAI DPPH. The TPC also gave negative and significant correlation with IC₅₀ tyrosinase (Table 3). This mean that when TPC increase, the AAI DPPH increase which showed high antioxidant activity. Increasing in TPC will correlate with decreasing in IC₅₀ of tyrosinase, which presented high tyrosinase inhibitory activity. This correlation result was in line with the results of the study which showed that ethanolic extract had the higher TPC than the DEA and DNs.

Identification of marker compound

Marker compound was used to control the consistency of crude drug, extract or product of the

extract. The results of the identification of compounds in the ethanol and DEAs by TLC co-chromatography (Figure 5) showed that there were spots of DE and DEA which had the same R_f value with quercitrin (0.38). The retention time of one peak in DE and DEA chromatograms coincided with the retention time on the quercitrin. Based on the HPLC chromatogram, quercitrin was dominant compound in DE and DEA. Then quercitrin was stated as marker compound in *P. campechiana* leaves extract. The HPLC chromatogram of DE and DEA of *P. campechiana* leaves and quercitrin standard was displayed in Figure 6. It confirmed the identification of marker compound of DE and DEA by TLC.




Figure 5: (a) TLC co-chromatography of DE and DEA with standard compounds: a. kaempferol, b. quercetin, c. apigenin, d. gallic acid, e. DEA, f. DE, g. chlorogenic acid, h. quercitrin, i. luteolin 7-O glycoside, j. rutin, monitoring under UV light λ 254 nm; (b) quercitrin structure




Figure 6: (a) Chromatogram HPLC of quercitrin; (b) DE; (c) DEA; (d) overlay quercitrin with DE; (e) overlay quercitrin with DEA

Marker compound content in *P. campechiana* leaves extract

Determination of marker compound content in DE and DEA were used one point method and the result showed in Table 4. The content of quercitrin in DEA was higher than in DE. Quercitrin is a compound that belongs to the flavonoid glycoside group. The benefit of quercitrin include being able to act as an

Table 4: Quercitrin content in ethanol and DEA of *Pouteria campechiana* leaves

Sample	Concentration	Retention time (min)	AUC	Quercitrin content
Quercitrin	500 µg/ml	6.998	15032928	-
DE	50000 µg/ml	6.930	2295223	0.153%
DEA	10000 µg/ml	6.972	10639291	3.539%

AUC: Area under curve, DE: Ethanol extract, DEA: Ethyl acetate extract.

antioxidant, antibacterial [21] and can protect the skin from exposure to UV B rays [22].

Conclusion

P. campechiana leaves extract had a potential as an antioxidant and tyrosinase inhibitor. DE of *P. campechiana* leaves had the highest potential for tyrosinase inhibitory and antioxidant activities. The highest TPC was in the DE and the highest TFC was in the DEA of *P. campechiana* leaves. TPC have a correlation with tyrosinase inhibitory and antioxidant activities. Quercitrin was chosen as a marker compound from *P. campechiana* leaves extract and the quercitrin content in the DEA was higher than in DE. Subsequent research will isolate compounds that can contribute to tyrosinase inhibitory and antioxidant activities.

Acknowledgment

The Author would like to thank to the Indonesia Endowment Fund for Education (LPDP) which has funded this research with contract number 2020032130798. Thanks are also extended to ITB and STFI who facilitated the instruments in carrying out this research, to Nurma Sabila who has helped in this research work for HPLC.

References

- Chunhakant S, Chaicharoenpong C. Antityrosinase, antioxidant, and cytotoxic activities of phytochemical constituents from *Manilkara zapota* L. Bark. *Molecules*. 2019;2798(24):1-19. <https://doi.org/10.3390/molecules24152798>
PMid:31370334
- Muddathir AM, Yamuchi K, Batubara I, Mohieldin EA, Mitsunaga T. Anti-tyrosinase, total phenolic content and antioxidant activity of selected Sundanese medicinal plant. *South Afr J Bot*. 2017;109:9-15. <http://dx.doi.org/10.1016/j.sajb.2016.12.013>
- Prota G. An introduction to melanin research. In: Prota, G. editor. *Melanins and Melanogenesis*. San Diego, CA, USA: Academic Press; 1992. p. 1-9.
- Ferrer S, Lopez AR, Carmon JG. Tyrosinase: A comprehensive

- reviewofitsmechanism. *BiochimBiophysActa*. 1995;1247(1):1-11. [https://doi.org/10.1016/0167-4838\(94\)00204-t](https://doi.org/10.1016/0167-4838(94)00204-t)
PMid:7873577
5. Kumar S, Pandey AK. Chemistry and biological activities of flavonoids: An overview. *Sci World J*. 2013;2013:162750. <https://doi.org/10.1155/2013/162750>
PMid:24470791
6. Batubara I, Darusman LK, Mitsunaga T, Rahminiwati M, Djauhari E. Potency of Indonesian medicinal plants as tyrosinase inhibitor and antioxidant agent. *J Biol Sci*. 2010;10(2):138-44. <https://doi.org/10.3923/jbs.2010.138.144>
7. Saraswaty V, Suparta NW, Setiyanto H, Rachmawati H, Adnyana IK. Transformation of melinjo seed micropowders into nanopowder enhances extractability of phenolic compounds and tyrosinase inhibitory activity. *Sains Malaysiana*. 2019;48(5):983-90. <https://dx.doi.org/10.17576/jsm-2019-4805-06>
8. Ma J, Yang H, Basile MJ, Kennelly EJ. Analysis of polyphenolic antioxidants from the fruit of three *Pouteria* species by selected ion monitoring liquid chromatography-mass spectrometry. *J Agric Food Chem*. 2004;52(19):5873-8. <https://doi.org/10.1021/jf049950k>
PMid:15366835
9. Hernandez CL, Villasenor IM, Joseph E, Tolliday N. Isolation and evaluation of antimutagenic activity of phenolic compounds from *Pouteria campechiana* Baehni. *Philippine J Sci*. 2008;137(1):1-10.
10. Pourmurad F, Hosseinimehr SJ, Shahabimajid N. Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. *Afr J Biotechnol*. 2006;5(11):1142-5.
11. Chang CC, Yang MH, Wen HM, Chen JC. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *J Food Drug Anal*. 2002;10(3):178-82. <http://doi.org/10.38212/2224-6614.2748>
12. Masuda T, Yamashita D, Takeda Y, Yonemori S. Screening for tyrosinase inhibitors among extract of seashore plants and identification of potent inhibitors from *Garcinia subelliptica*. *Biosci Biotechnol Biochem*. 2005;69(1):197-201. <https://doi.org/10.1271/bbb.69.197>
PMid:15665485
13. Blois MS. Antioxidant determinations by the use of a stable free radical. *Nature* 1958;181:1199-200.
14. Scherer R, Godoy HT. Antioxidant activity index (AAI) by the 2,2-diphenyl-1-picrylhydrazyl method. *Food Chem*. 2009;112:654-8. <https://doi.org/10.1016/j.foodchem.2008.06.026>
15. Tanase C, Cosarca S, Muntean DL. A critical review of phenolic compounds extracted from the bark of woody vascular plants and their potential biology activity. *Molecules*. 2019;24(1182):1182. <http://doi:10.3390/molecules24061182>
PMid:30917556
16. Treml J, and Smejkal K. Flavonoids as potent scavengers of hydroxyl radicals. *Compr Rev Food Sci Food Saf*. 2016;15(1):720-38. <https://doi.org/10.1111/1541-4337.12204>
17. Chen CY, Lin LC, Yang WF, Bordon J, Wang HMD. An update organic classification of tyrosinase. *Curr Org Chem*. 2015;19(1):4-18. <http://doi:10.2174/1385272819666141107224800036>
18. Kim YJ, Uyama H. Review: Tyrosinase inhibitors from natural and synthetic sources: Structure, inhibition mechanism and perspective. *Cell Mol Life Sci*. 2005;62(15):1707-23. <http://doi:10.1007/s00018-005-5054-y>
PMid:15968468
19. Rao GV, Sahoo MR, Madhavi MS, Mukhopadhyay T. Phytoconstituents from the leaves and seeds of *Manilkara zapota* Linn. *Pharm Lett*. 2014;6(2):69-73.
20. Baky MH, Kamal AK, Elgindi MR, Haggag EG. A review on phenolic compounds from family sapotaceae. *J Pharmacogn Phytochem*. 2016;5(2):280-7.
21. Hardiyanti R, Marpaung L, Andiyana IK, Simanjuntak P. Isolation of quercitrin from *Dendrophthoe pentandra* (L.) and its antioxidant and antibacterial activities. *Rasayan J Chem*. 2019;12(4):1822-7. <http://dx.doi.org/10.31788/RJC.2019.1235353>
22. Yin Y, Li W, Son Y, Sun L, Lu J, Kim D, et al. Quercitrin protects skin from UVB-induced oxidative damage. *Toxicol Appl Pharmacol*. 2013;269(2):88-99. <http://dx.doi.org/10.1016/j.taap.2013.03.015>
PMid:23545178