

Berkas Publikasi Sebagai Co Author : Pharmacology and Clinical Pharmacy Research

Comparison of Anti-Inflammatory Activity between Fermented and Dried Breadfruit Leaves Extract (*Artocarpus Altilis*)

Hesti Riasari, Sani N Fitriansyah, Oktamuman Putra

Sekolah Tinggi Farmasi Indonesia

1. Halaman sampul
2. Dewan Redaksi/Editorial Board
3. Daftar isi
4. Jurnal Published

1. Halaman Sampul Jurnal

Pharmacology and Clinical Pharmacy Research

ISSN : 2527-7332
e-ISSN : 2614-0020

Accredited SINTA 2 by The Ministry of Research and Technology of Indonesia (No.204/E/KPT/2022).

HOME ABOUT LOGIN REGISTER CATEGORIES SEARCH CURRENT ARCHIVES

Home > Vol 9, No 1 (2024)

Pharmacology and Clinical Pharmacy Research

Pharmacology and Clinical Pharmacy Research (PCPR) is an international, peer-reviewed, scientific journal that focus on pharmacology and clinical pharmacy. PCPR is published three times a year (April, August, and December) to provide a forum for pharmacologists, clinical pharmacists, and other researchers to share the current update on pharmacology and best practice in clinical pharmacy.

PHARMACOLOGY AND CLINICAL PHARMACY RESEARCH has been accredited SINTA 2 by The Ministry of Research and Technology of Indonesia (No. 204/E/KPT/2022). PCPR is already indexed in DOAJ, Cross-Ref, Google Scholar, Index Copernicus, and Garuda-Ristekdikti. All articles published by PCPR are made immediately available worldwide under an open access license.

PCPR welcomes submission of original research articles, review articles, and case reports. All submitted manuscripts will go through the double-blind peer review before being granted with acceptance for publication. It is published by Faculty of Pharmacy, Universitas Padjadjaran, Indonesia. Scope of the journal includes:

- Experimental Pharmacology
- Clinical Pharmacology
- Molecular Pharmacology
- Toxicology
- Pharmacokinetics

ONLINE SUBMISSIONS
FOCUS AND SCOPE
AUTHOR GUIDELINES
PUBLICATION ETHICS
EDITORIAL TEAM
PEER-REVIEWERS
SUBSCRIPTION/ORDER
INDEXING
CHAT VIA WHATSAPP

INDEXED BY 



2. Editorial Board

Pharmacology and Clinical Pharmacy Research

ISSN : 2527-7332
e-ISSN : 2614-0020

Accredited SINTA 2 by The Ministry of Research and Technology of Indonesia (No.204/E/KPT/2022).

[HOME](#) [ABOUT](#) [LOGIN](#) [REGISTER](#) [CATEGORIES](#) [SEARCH](#) [CURRENT](#) [ARCHIVES](#)

[Home](#) > [About the Journal](#) > [Editorial Team](#)

Editorial Team

Editor in Chief

Jutti Levita, Faculty of Pharmacy, Universitas Padjadjaran, Indonesia

Executive Editor

Rizky Abdulah, Faculty of Pharmacy, Universitas Padjadjaran, Indonesia

Editorial Board

Debabrata Banerjee, Department of Pharmacology, Rutgers University, United States
Dyah A. Perwitasari, Faculty of Pharmacy, Ahmad Dahlan University, Indonesia
Edy Meiyanto, Faculty of Pharmacy, Gajah Mada University, Indonesia
Hiroshi Koyama, Department of Public Health, Gunma University, Japan
Ivan Surya Pradipta, University of Groningen, Netherlands
Prof. Dr. apt. Keri Lestari, M.Si., Faculty of Pharmacy, Universitas Padjadjaran, Indonesia
Henk-Jan Guchelaar, Department of Clinical Pharmacy and Toxicology, Leiden University, Netherlands
John K. Baird, Eijkman-Oxford Clinical Research Unit, Oxford University, United Kingdom
Lyn T. Le, School of Biotechnology, Vietnam National University, Viet Nam
Maarten J. Postma, Department of Pharmacy, University of Groningen, Netherlands
Syed AS. Sulaiman, School of Pharmaceutical Sciences, Universiti Sains Malaysia, Malaysia
Supat Jiranusornkul, Faculty of Pharmacy, Chiang Mai University, Thailand

[ONLINE SUBMISSIONS](#)

[FOCUS AND SCOPE](#)

[AUTHOR GUIDELINES](#)

[PUBLICATION ETHICS](#)

[EDITORIAL TEAM](#)

[PEER-REVIEWERS](#)

[SUBSCRIPTION/ORDER](#)

[INDEXING](#)

[CHAT VIA WHATSAPP](#)

INDEXED BY



USER

[\[User Information\]](#)

3. Daftar Isi

Vol 3, No 3

DOI: <https://doi.org/10.15416/pcpr.v3i3>

Table of Contents

Original Article

Formulation and Evaluation of Anti-Acne Lotion Containing Red Ginger (<i>Zingiber officinale</i> Roscoe) Essential Oil <i>Dwi Indriati, Ike Y. Wiendarlina, Annisa S. Carolina</i>	PDF 61-65
Prescription Pattern and Costs of Angiotensin II Receptor Blockers in Hypertensive Outpatients <i>Nurmajnah Nurmajnah, Nur Anisyah</i>	PDF 66-71
Comparison of Anti-Inflammatory Activity between Fermented and Dried Breadfruit Leaves Extract <i>Hesti Riasari, Sani N. Fitrianyah, Oktamuman Putra</i>	PDF 72-75
Should Cannabis as Medicine be Specifically Regulated? <i>Gunawan Widjaja</i>	PDF 76-79
In Vitro Inhibition of α -Amylase and α -Glucosidase by Different Parts of Amla Plant (<i>Phyllanthus emblica</i> L) <i>Novi I. Fauzi, Maria Ulfah, Syarif Hamdani, Saeful Hidayat</i>	PDF 80-84
Authors and Keywords Index <i>Editorial PCPR</i>	PDF

FOCUS AND SCOPE

AUTHOR GUIDELINES

PUBLICATION ETHICS

EDITORIAL TEAM

PEER-REVIEWERS

SUBSCRIPTION/ORDER

INDEXING

CHAT VIA WHATSAPP

INDEXED BY



USER

Username

Password

Remember me

MANUSCRIPT TEMPLATE

Comparison of Anti-Inflammatory Activity between Fermented and Dried Breadfruit Leaves Extract (*Artocarpus Altilis*)

Hesti Riasari, Sani N.Fitriansyah, Oktamuman Putra
Department of Biology Pharmacy, Indonesia School of Pharmacy, Bandung,
West Java, Indonesia

Abstract

Breadfruit leaves (*Artocarpus altilis*) have been empirically used to treat inflammation. Biosynthesis process such as fermentation can possibly changes the composition of secondary metabolites in breadfruit leaves. The aim of this study was to compare the anti-inflammatory activity between fermented and natural-dried breadfruit leaves extract. Fermented and dried breadfruit leaves were extracted with soxhlet device using methanol solvent. The anti-inflammatory activity of both extracts were evaluated using carrageenan (1% w/v) foot edema model of inflammation in male Wistar rats. The edema was measured with plethysmometer every 1 hour for 6 hours. The results showed that both extracts exhibited anti-inflammatory activity. 50 mg and 100 mg fermented leaves extracts exhibited 37.17% and 71.82% inhibition, while the same concentrations of dried extract exhibited 39.50% and 57.72%, inhibition respectively. In conclusion, the best anti-inflammatory activity was shown by fermented breadfruit leaves extract.

Keywords: *Artocarpus altilis*, anti-inflammatory, breadfruit, fermentation

Introduction

Indonesia has the highest biodiversity in the world. Of 30 thousands of plants species, more than 1000 species have been used for medical treatment purpose. Breadfruit plant (*Artocarpus altilis*) is one of the plants that has been empirically used to treat inflammation, skin diseases, and liver cirrhosis.^{1,2}

Previous study showed that breadfruit leaves (60 mg/kg) had potent anti-inflammatory effect.³ A study conducted by Riasari *et al* showed that cytostenone which was successfully isolated from the breadfruit

leaves is considered responsible for its anti-inflammatory activity.³⁻⁵ Biosynthesis process, such as fermentation can possibly changes the composition of secondary metabolites in breadfruit leaves.⁶

Currently, there is limited information whether the fermentation process could enhance anti-inflammatory activity of the breadfruit leaves. Therefore, this study was conducted to compare anti-inflammatory activity between fermented and dried breadfruit leaves extract.

Corresponding author: Hesti Riasari. Department of Biology Pharmacy, Indonesia School of Pharmacy. Bandung, West Java, Indonesia. Email: hmm_riasari@yahoo.com

Received: 22 October 2018. **Revised:** 14 November 2018. **Published:** 19 December 2018.

Table 1. % Inhibition of Breadfruit Leaves Extract on Inflammation

Dosage Treatment	T1(%)	T2(%)	T3(%)	T4(%)	T5(%)	T6(%)
Positive control	57.93	63.82	43.05	47.11	59.60	77.56
50 mg fermented leaves extract	41.11	27.51	27.51	32.69	37.17	37.17
100 mg fermented leaves extract	47.19	67.12	67.12	69.83	71.82	71.82
50 mg dried leaves extract	27.08	10.26	30.21	35.19	39.50	39.50
100 mg dried leaves extract	34.37	41.67	50.24	54.16	57.72	57.72

Methods

Plant determination was conducted at Herbarium, School of Biotechnology Science, Bandung Institute of Technology. Instruments used in this study included soxhlet extraction device, rotary evaporator, and plethysmometers. Materials used in this study included natural-dried breadfruit leaves, fresh breadfruit leaves, methanol solvent, male Wistar rats, 0.9% sterile NaCl, carrageenan, PGA, sodium diclofenac and distilled water. Fresh breadfruit leaves were washed, dried, and fermented for 5 days. After the fermentation process was complete, the leaves were dried at room temperature.

Fermented and natural-dried breadfruit leaves were assayed to determine ash content, percentage of ethanol soluble compounds, and percentage of water soluble compounds. Extraction was performed using soxhlet device at 60 °C. A total of 163.31 g of fermented breadfruit leaves were extracted with 1.8 l of methanol solvent. Extract of dried breadfruit leaves was obtained by extracting 150 g of leaves with 1.8 l of methanol solvent. During extraction, the leaves were wrapped with filter paper and placed in the soxhlet device. The methanol solvent was placed into a flask and heated at approximately 60 °C. The solvent was evaporated and underwent condensation in the cooling device, before extracting the plant. The liquid extract was then concentrated using a rotary evaporator

to obtain thick extracts. Phytochemical screening was performed to determine the composition of secondary metabolites in both extracts. The compounds tested included alkaloids, flavonoids, tannins, phenolics, triterpenoids, steroids, quinones, monoterpenes, sesquiterpenes, and saponins.

The anti-inflammatory activity of both extracts were evaluated using carrageenan (1% w/v) foot edema model of inflammation in the male Wistar rats. Carrageenan was used as inducer of inflammation since it has more sensitive response to drugs.^{7,8} The edema was measured with plethysmometer every 1 hour for 6 hours. Analysis of variance (ANOVA) was conducted. $P < 0.05$ defined statistical significance. The activities of the extracts were compared with standard drug, sodium diclofenac.

Results and Discussion

Results of the plant determination showed that plant used in this study was *Artocarpus altilis* (Parkinson) Fosberg. Phytochemical screening showed that natural-dried breadfruit leaves contained flavonoids, steroids, triterpenes, monoterpenes, sesquiterpenes and phenols, while fermented breadfruit leaves contained flavonoids, steroids, monoterpenes, sesquiterpenoids, phenols, quinones and tannins. Our study showed that fermentation process resulted in

different secondary metabolites. Ash content in natural-dried breadfruit leaves was 37%, while in fermented breadfruit leaves was 20.50%. Determination of ash content is important to examine content of mineral and anorganic elements. Water soluble content in dried breadfruit leaves was 2%, while in the fermented leaves was 21.9%. In dried breadfruit leaves, ethanol soluble content was 2.2%, while in the fermented leaves was 4.95%. These tests were important to provide information related to dissolved compounds and to select appropriate solvent for extraction. This finding showed that fermented breadfruit leaves contained more polar compounds.

Soxhlet extraction was performed because secondary metabolites in breadfruit leaves were heat-resistant. Methanol was selected because it had better polarity compared to other solvents, thus it could extract more secondary metabolites. We obtained 8.91 g (5.94%) thick extract of natural-dried breadfruit leaves and 20.94 g (12.82%) thick extract of fermented breadfruit leaves.

% inhibition against inflammation can be seen in Table 1. We found that strongest inhibition was shown by 100 mg of fermented leaves extract (71.82%). The lowest inhibition was shown by 50 mg of fermented leaves extract (37.17%) and 50 mg of dried leaves extract (39.50%). Based on statistical analysis, 100 mg of fermented leaves extract had the best anti-inflammatory activity compared to other extracts. Geranyl flavonoids and its derivatives isolated from the breadfruit leaves were considered responsible for its anti-inflammatory activity.^{9,10}

Conclusion

The strongest anti-inflammatory activity was exhibited by fermented breadfruit leaves extract.

Acknowledgement

None declared.

Funding

None.

Conflict of interest

None declared.

Reference

1. Jagtap UB, Bapat VA. Artocarpus: a review of its traditional uses, phytochemistry and pharmacology. *Journal of Ethnopharmacology*. 2010;129(2):142–166.
2. Ramalingum N, Mahomoodally MF. The therapeutic potential of medicinal foods. *Advances in Pharmacological Science*. 2014;2014:354264.
3. Lee CW, Ko HH, Chai CY, Chen WT. Effect of *Artocarpus communis* extract on UVB irradiation-induced oxidative stress and inflammation in hairless mice. *International Journal of Molecular Science*. 2013;14(2):3860-3873.
4. Sairam S, Urooj A. Safety evaluation of *Artocarpus altilis* as pharmaceutical agent in Wistar rats. *Journal of Toxicology*. 2014;2014:980404.
5. Riasari H, Sukrasno, Komar R. Metabolite profile of various development bread fruit leaves (*Artocarpus altilis*. parkinson fosberg) and the identification of their major components. *International Journal of Pharmaceutical Science and Research*. 2015;6(5): 2170-2177.
6. Yang L, Wen KS, Ruan X, Zhao YX, Wei F, Wang Q. Response of plant secondary metabolites to environmental factors. *Molecules*. 2018;23(4):762.
7. Kumar T, Jain V. Anti-nociceptive and anti-inflammatory activities of *Bridelia retusa* methanolic fruit extract in experimental animals. *Scientific World*

- Journal*. 2014;2014:890151.
8. Mansouri MT, Hemmati AA, Naghizadeh B, Mard SA, Rezaie A, Ghorbanzadeh B. A study of the mechanisms underlying the anti-inflammatory effect of ellagic acid in carrageenan-induced paw edema in rats. *Indian Journal of Pharmacology*. 2015;47(3):292-8.
 9. Hsu CL, Chang FR, Tseng PY. Geranyl flavonoid derivatives from the fresh leaves of *Artocarpus communis* and their anti-inflammatory activity. *Planta Medica*. 2012;78(10):995-1001.
 10. Lin JA, Chen HC, Yen GC. The preventive role of breadfruit against inflammation-associated epithelial carcinogenesis in mice. *Molecular Nutrition and Food Research*. 2014;58:206–210.